

Relativistic Particle Acceleration in a Folded Current Sheet

S. Zenitani and M. Hoshino

*Department of Earth and Planetary Science, University of Tokyo, 7-3-1, Hongo, Bunkyo,
Tokyo, 113-0033 Japan; zenitani@eps.s.u-tokyo.ac.jp*

ABSTRACT

Two-dimensional particle simulations of a relativistic Harris current sheet of pair ($e^\pm$) plasmas have demonstrated that the system is unstable to the relativistic drift kink instability and that a new kind of acceleration process takes place in the deformed current sheet. This process contributes to the generation of non-thermal particles and contributes to the fast magnetic dissipation in the current sheet structure. The acceleration mechanism and a brief comparison with relativistic magnetic reconnection are presented.

Subject headings: acceleration of particles — instabilities; plasmas — relativity; magnetic fields — pulsars(individual: Crab Pulsar)

1. Introduction

A current sheet structure with reversed magnetic fields can be seen everywhere in the universe. Among the magnetic field dissipation processes in the current sheet, magnetic reconnection may be the most popular one. Breaking the topology of magnetic field lines, it releases the magnetic energy into particles' kinetic energy in a relatively short time-scale (several to several tens of an Alfvén transit time $\tau_A = \lambda/V_A$, where λ is the half thickness of the current sheet and V_A is the typical Alfvén speed). It plays an essential role in solar terrestrial sites such as the Earth's magnetotail (Dungey 1961) and source regions of the solar flare (Parker 1957). Moreover, it is expected in astrophysical sites such as active galactic nuclei (Lesch & Birk 1998; Schopper et al. 1998) and pulsar winds, especially their well-studied example of the Crab Nebula (Coroniti 1990). In case of the Crab Nebula, energy conversion in a relativistic wind of $e^\pm$ plasmas has been a long-standing problem (the so-called the “ σ -problem”, where σ is the ratio of the Poynting flux energy to the particle kinetic flux) (Kennel & Coroniti 1984; Coroniti 1990; Lyubarsky & Kirk 2001; Kirk & Skjæraasen 2003). The wind is originally Poynting-dominated ($\sigma \sim 10^4$) close to the neutron star (Arons 1979); but it is kinetic-dominated near the termination shock (σ is far less than

the unity; $\sigma \sim 10^{-3}$ by Kennel & Coroniti (1984) and $\sigma \sim 10^{-1}$ by Mori et al. (2004)). Recently, relativistic magnetic reconnection in the striped current sheets has been discussed as the most possible dissipation mechanism (Coroniti 1990; Lyubarsky & Kirk 2001; Kirk & Skjæraasen 2003). The properties of relativistic reconnection in $e^\pm$ plasmas have been investigated by an analytical study (Blackman & Field 1994) and by computer simulations (Zenitani & Hoshino 2001; Jaroschek et al. 2004). One of the most important features of relativistic reconnection is the generation of non-thermal particles due to direct acceleration by dc electric field around the X -type region (Zenitani & Hoshino 2001). However, the above discussion is based on a two-dimensional picture of reconnection. Instabilities in a cross-field plane are also of critical importance to understanding a realistic current sheet problem that includes reconnection. Does reconnection grow faster than cross-field instabilities? Is reconnection the fastest dissipation process? In the case of non-relativistic ion-electron plasmas, the following instabilities have been discussed: the lower hybrid drift instability (LHDI) (Krall 1971; Davidson et al. 1977), the Kelvin-Helmholtz instability (KHI) (Yoon & Drake 1996; Shinohara et al. 2001), and the drift kink instability (DKI) (Zhu & Winglee 1996; Pritchett et al. 1996; Daughton 1998). They lead to plasma heating and particle acceleration as well as the triggering of reconnection. The purpose of this Letter is to search for the basic properties of such cross-field instabilities and of the non-thermal particle generation in a relativistic $e^\pm$ current sheet. In this Letter, we report our 2D simulations in a cross-field plane of a relativistic $e^\pm$ current sheet and we report a new acceleration process related to the relativistic drift kink instability (RDKI) (Zenitani & Hoshino 2003).

2. Simulation

The simulation code used here is a three-dimensional particle-in-cell code. The system size is composed of 1 (X) $\times$ 256 (Y) $\times$ 512 (Z) grids. We consider periodic boundaries in the X , Y directions, and we periodically set two simulation boxes in the Z direction. The typical scale of the current sheet λ is set to 10 grids, so that the Y boundaries are located at $y = \pm 12.8\lambda$, and the Z boundaries are located at $z = \pm 12.8\lambda$. We take the relativistic Harris model as the initial current sheet. The magnetic field, the density of plasmas and their distribution functions are described by $\mathbf{B} = B_0 \tanh(z/\lambda) \hat{\mathbf{x}}$, $n(z) = n_0 \cosh^{-2}(z/\lambda)$ and $f_\pm \propto n(z) \exp[-\Gamma_\beta \{\varepsilon - \beta_\pm u_y\}/T]$, respectively. In the above equations, B_0 is the magnitude of magnetic field in the lobe (background region), n_0 is the number density of plasmas in the current sheet, $\beta_\pm = v_\pm/c$ are the drift velocities for each species; $\beta_+ = +\beta$ for positrons, $\beta_- = -\beta$ for electrons, Γ_β is the Lorentz factor for β [$\Gamma_\beta = [1 - \beta^2]^{-1/2}$], ε is the particle energy, $\mathbf{u}$ is the relativistic four velocity of $\mathbf{u} = [1 - (v/c)^2]^{-1/2} \cdot \mathbf{v}$ and T is the temperature. In the first simulation, we set $T = mc^2$ and $\beta = 0.3$. We set no driving force to excite

instabilities, so that instabilities arise from thermal noise. We set no background plasmas. The total energy is conserved within an error of 0.3% throughout the simulation run.

3. Results

Snapshots at three characteristic stages of the simulation are presented in Figure 1. The three left panels in Figure 1 show color contours of the plasma density at (a) $t/\tau_c = 46.0$, (b) 64.0 and (c) 82.0, where time is normalized by the light transit time $\tau_c = \lambda/c$. The three middle panels show color contours of the electric field E_y at the corresponding time; the white lines show contours of B_x . The three panels show the particles' energy spectra for the whole simulation. The horizontal axis shows the particles' energy normalized by mc^2 .

As shown in the top three panels in Fig. 1, we observe a kink-type modulation of the current sheet. We confirmed that this is a signature of the RDKI, a relativistic extension of the DKI, which is a long-wavelength, current-driven instability in a thin current sheet. In the top middle panel, one can see the typical structure of E_y components of the polarization of electric fields of RDKI. The sign of E_y is positive in the yellow regions, while it is negative in the blue regions. Two types of regions appear alternatively along the current sheet and they are anti-symmetric with the neutral plane ($z = 0$). Note that the $\mathbf{E} \times \mathbf{B}$ direction using the lobe magnetic fields and the observed sign of E_y is consistent with the Z -displacement of the plasma bulk motion. The symmetric E_z components are also observed and their values are a little less than or comparable to those of E_y . The E_x components are negligible. Since the system is uniform in the X direction, there is no electrostatic E_x fields. After $t/\tau_c \sim 50$, the instability turns into its non linear stage. The three middle panels in Fig. 1 are snapshots of the current sheet at $t/\tau_c = 64.0$. The current sheet is strongly folded within a zone of $-4 < z/\lambda < 4$. Due to the Z -displacement by RDKI, red (originally yellow) regions, where $E_y > 0$, are placed in a row around the neutral plane ($z = 0$) in the midmost panel of Fig. 1. The total kinetic energy in the system has increased by 30% at this time. Importantly, we observe a clear sign of the particle acceleration in a high-energy tail in the energy spectrum of the middle-right panel in Fig. 1. We will discuss the acceleration process later. The kinked-shape of the current sheet is not stable, because fragments of the anti-parallel currents ($\pm J_z$) pull each other. As they start to collide, the system turns into the “mixed” stage. The snapshots at this stage are presented in the bottom three panels in Fig. 1 at $t/\tau_c = 82.0$. Due to the enhanced diffusion by sheet collisions, the total kinetic energy has increased by 170% at this time. In the energy spectrum of the bottom-right panel, one can recognize both a remnant of the non-thermal tail and global plasma heating. Finally, the mixed current sheet slowly evolves into the broadened current sheet, which is 3-4 times

thicker than the initial state. The saturated level of the total kinetic energy is 330%-340% at $t/\tau_c = 200$ or later. The spectrum looks unchanged from the mixed stage; one can still observe the non thermal tail.

4. Discussion

Now, we again take a look at the instability during its linear stage. The observed growth rate (ω_i/Ω_c) is plotted as a function of $k_y\lambda$ in Fig. 2, where k_y is the wavenumber and Ω_c is the gyro frequency $\Omega_c = \omega_c/\gamma = (eB)/(\gamma mc^2)$. The growth rate and wave length of the most dominant mode of RDKI are $\omega_i/\Omega_c \sim 0.035$ and $k_y\lambda \sim 1.0$, respectively. We have also calculated eigen functions and their growth rates of the RDKI, by linearizing relativistic two-fluid equations. The theoretical growth rates for the present case are shown as a solid line in Fig. 2. The simulation results are consistent with them except for shorter wavelength of $k_y\lambda > 1$, where the instability is suppressed by kinetic effects. We observe a slight signature of the relativistic drift sausage instability (RDSI) along with RDKI. However, RDSI is less influential on the current sheet deformation and particle acceleration. Note that no velocity-shear-driven mode such as KHI is excited, and that is because there is no velocity shear between drifting plasmas in the current sheet and empty background plasmas. Also note that LHDI is not excited in an $e^\pm$ case, because there is no delay of a positron's response to an electron's.

Next, we discuss an acceleration mechanism that is found at the non linear stage of RDKI. By analyzing trajectories of highly accelerated particles, we found that the main site of particle acceleration is the central channel around $z \sim 0$, where the current sheet was originally located. For simple discussion, we call this channel the “acceleration channel (AC)”. We illustrate the acceleration mechanism in schematic views in Fig. 3. In the upper view, the structure of the polarization of E_y fields at the linear stage are illustrated. They are anti symmetric with respect to Z and both positive and negative regions are alternatively located along the sheet. Passing through their meandering orbits, some particles can resonate with the E_y fields. Some of them gain their energy and others lose their energy. At the non-linear stage, due to the Z -displacement of the current sheet, the positive E_y regions are now located around the AC. Inside the positive E_y regions, typical values of the fields are $|B_x| \approx 0.6 - 0.7B_0$ and $E_y \approx 0.2 - 0.3B_0$. Contrary to the reconnection (Zenitani & Hoshino 2001; Jaroschek et al. 2004) in which $|E|/|B| > 1$ in an acceleration site, the value of $|E|/|B|$ is less than unity inside the positive E_y regions, and so low-energy particles travel into the $E \times B$ direction. However, fewer populations of high-energy particles can travel across the positive E_y regions, because their Larmor radii r_L are larger than the half-wavelength of

RDKI (π/k_y). They gain their energy from the electric field E_y , cross the current sheet and move into the neighboring positive E_y regions. Since r_L becomes larger as particles gain more energy ($r_L \propto \varepsilon^{1/2}$ in non-relativistic case and $r_L \propto \varepsilon$ in relativistic case), accelerated particles usually satisfies the crossing condition of $r_L > \pi/k_y$ in the following positive E_y regions. In this way, they are successively accelerated in the AC, passing through multiple positive E_y regions into the $\pm Y$ direction. The above condition can be interpreted as the “trapping condition” inside the AC. Higher accelerated particles tend to stay in the AC, because larger r_L prevents them from escaping. This is why the non-thermal tail is enhanced in the energy spectrum. We think this type of enhanced acceleration can be found in any kink-type instability in a thin current sheet, as long as its wavelength is comparable to a particle’s Larmor radius. For example, it may be applicable to acceleration of ions, when a thin current sheet of ion-electron plasmas is modulated by KHI. The folded structure evolves into the mixed stage when its Z -displacement (Δz) is nearly the same as the half-wavelength of RDKI ($\Delta z \sim \pi/k_y \sim \pi\lambda$). Assuming that v_z is typically $V_{E \times B} \sim 0.3c$ in the positive E_y regions, the time scale of the non-linear stage (τ_N) is approximated by $\tau_N \sim \pi\lambda/v_z \sim 5-10\tau_c$.

After the system turns into the mixed stage, a few number of high-energy particles and a lot of low-energy particles in the current sheet interact with the positive E_y regions. Then the magnetic fields are dissipated in a time-scale of tens of τ_c . Roughly speaking, the magnetic field energy is consumed by Joule heating of the E_y fields and the total current $\bar{I}_y$ in the broadened current sheet. The magnetic energy stored between the folded curves is approximated by $(\pi/k_y)B_0^2/8\pi$. We assume that the average value of the electric fields $\bar{E}_y \sim 0.5E_y = 0.5 \times (0.2 - 0.3)B_0$, where the factor 0.5 represents the fact that positive E_y is not uniform in the AC. The total current $\bar{I}_y$ is represented by the zeroth order current $\bar{I}_y \sim \lambda J_0 = cB_0/4\pi$. Thus the dissipation time scale (τ_D) is approximated by $\tau_D \sim (\pi/k_y)(B_0^2/8\pi)(\lambda J_0 \bar{E}_y)^{-1} \sim 10 - 20\tau_c$.

Next, we briefly compare RDKI and magnetic reconnection (Zenitani & Hoshino 2001). In Table 1, the parameters for several simulation runs of RDKI and for relevant runs of reconnection are presented. Throughout the simulation runs, we calculate the ratio of the non-thermal kinetic energy to the total kinetic energy (K_{nonth}/K) and their maximum values in time are presented in Table 1. In case of run D3, K_{nonth} is not resolved by our method. The maximum energies (ε_{max}) are also presented in Table 1. In the case of RDKI, the highest energy particles run in the AC into the $\pm Y$ direction with the light speed during the non-thermal stage and the mixed stage. Their maximum energies are estimated by $\varepsilon_{est} \sim \varepsilon_{max0} + ec\bar{E}_y(\tau_N + \tau_D) \sim \varepsilon_{max0} + 3\omega_c\tau_c mc^2$ and the estimated values agree with the simulation results. In three cases of reconnection, the evolution seems to be restricted by the periodic system size of L_x , because two outflow jets travel into the $\pm X$ direction. Due to this restriction, the upper limits of particle energy are estimated by $\varepsilon_{est} = eB_0L_x/2$ in Table 1

in reconnection cases. However, the obtained maximum energies (ε_{max}) by reconnection are far larger than the maximum energies by RDKI. From the viewpoint of field dissipation (e.g. striped pulsar winds), RDKI can dissipate more magnetic energy. The linear growth rates of the energy conversion ($\tau_c d \ln(\Delta K)/dt$) are presented in Table 1. Generally speaking, RDKI grows *faster* in a relativistic current sheet ($T/mc^2 \geq 1$) and it converts 2.5 – 3.5 times as much energy as the initial kinetic energy in the current sheet in a time scale of τ_D . Moreover, RDKI spontaneously occurs everywhere in the X direction, while single reconnection requires a wide spatial range in the X direction in order for the outflow jets to escape. Therefore, even in runs R3 and D3, in which RDKI grows slower, reconnection takes twice as long to dissipate the same amount of the magnetic energy as RDKI. On the contrary, reconnection is the more favored candidate as the origin of non-thermal particles. It produces more non-thermal energy of K_{nonth} than RDKI as a result of the high ratio of K_{nonth}/K .

Our initial results of a three-dimensional simulation show that RDKI occurs almost uniformly in the current sheet. Also, Jaroschek et al. (2004) have presented their three-dimensional results of triggered reconnection. They have referred to the importance of RDKI in the three-dimensional case and a signature of a two-dimensional-like RDKI is observed outside the reconnection region. In case of multiple current sheets structure, such as striped pulsar winds, there are several possibilities of enhanced field dissipation. If the distance between two currents is less than the critical distance of $2\lambda_c$, the two current sheets collide with each other and 90% or more of the magnetic energy will be dissipated into the particle energy. In typical cases, $\lambda_c \sim 7-8$, but its dependency on T or β should be further investigated. We also note that the secondary RDKI takes place in the broadened current sheet in some occasions.

Let us summarize this Letter. First, we investigate the RDKI in a thin $e^\pm$ current sheet. Second, we find a new acceleration process that uses a folded structure of a current sheet. The unique point of this process is that AC electric fields evolve into a DC acceleration channel. Third, due to the enhanced acceleration, a non-thermal particles are produced. Fourth, comparison with reconnection shows that RDKI is more favorable to dissipate the magnetic energy in the current sheet structure. We believe that RDKI provides a crucial clue to the energy dissipation problem in the current sheets in the universe, including striped pulsar winds.

This work was supported by the facilitates of JAXA and the Solar-Terrestrial Environment Laboratory, Nagoya University.

REFERENCES

- J. Arons 1979, *Space Sci. Rev.*, 24, 437
- E. G. Blackman and G. B. Field 1994, *Phys. Rev. Lett.*, 72, 494
- F. V. Coroniti 1990, *ApJ*, 349, 538
- W. Daughton 1998, *J. Geophys. Res.*, 103, 29429
- R. C. Davidson et al. 1977, *Phys. Fluids*, 20, 301
- J. W. Dungey 1961, *Phys. Rev. Lett.*, 6, 47
- C. H. Jaroschek, R. A. Treumann, H. Lesch and M. Scholer 2004, *Phys. Plasmas*, 11, 1151
- C. F. Kennel and F. V. Coroniti 1984, *ApJ*, 283, 694
- J. G. Kirk and O. Skjæraasen 2003, *ApJ*, 591, 366
- N. A. Krall 1971, *Phys. Rev. A*, 4, 2094
- H. Lesch and G. T. Birk 1998, *ApJ*, 499, 167
- Y. Lyubarsky and J. G. Kirk 2001, *ApJ*, 547, 437
- K. Mori, D. N. Burrows, J. J. Hester, G. G. Pavlov, S. Shibata and H. Tsunemi 2004, *ApJ*, 609, 186
- E. N. Parker 1957, *J. Geophys. Res.*, 62, 509
- P. L. Pritchett, F. V. Coroniti and V. K. Decyk 1996, *J. Geophys. Res.*, 101, 27413
- R. Schopper, H. Lesch and G. T. Birk 1998, *A&A*, 335, 26
- I. Shinohara, H. Suzuki, M. Fujimoto and M. Hoshino 2001, *Phys. Rev. Lett.*, 8709, 5001
- P. Yoon and J. F. Drake 1996, *J. Geophys. Res.*, 101, 27327
- S. Zenitani and M. Hoshino 2001, *ApJ*, 562, L63
- S. Zenitani and M. Hoshino 2003, *Proc. of the 28th ICRC*
- Z. Zhu and R. M. Winglee 1996, *J. Geophys. Res.*, 101, 4885




Fig. 1.— Left three panels: Color contours of the plasma density at $t/\tau_c = 46.0$, 64.0 and 82.0 . Middle three panels: E_y at the corresponding time; the white lines shows contours of B_x . Right three panels: Energy spectra.




Fig. 2.— Plot of the observed growth rates (ω_i / Ω_c) as a function of the normalized wavenumber ($k_y \lambda$). The solid line shows the growth rate predicted by relativistic two-fluid theory.




Fig. 3.— Schematic view of the acceleration mechanism. Once the current sheet is folded at the non linear stage, regions with the positive E_y sign are located in a central acceleration channel.

Run	T	L_x	L_y	L_z	K_{nonth}/K	$\tau_c d \ln(\Delta K)/dt$	ε_{max}	ε_{est}
D1	16	1.0	25.6	51.2	4.7×10^{-2}	2.3×10^{-1}	7.0×10^2	7.1×10^2
D2	1	1.0	25.6	51.2	7.2×10^{-2}	2.3×10^{-1}	4.8×10^1	4.2×10^1
D3	1/16	1.0	25.6	51.2	-	1.9×10^{-2}	4.0	3.0
R1	16	102.4	1.0	51.2	$> 2.5 \times 10^{-1}$	6.6×10^{-2}	$> 3.6 \times 10^3$	5.6×10^3
R2	1	102.4	1.0	51.2	$> 2.6 \times 10^{-1}$	6.5×10^{-2}	$> 1.2 \times 10^2$	3.5×10^2
R3	1/16	102.4	1.0	51.2	$> 1.1 \times 10^{-1}$	7.1×10^{-2}	> 5.3	2.2×10^1

Table 1: Three runs for RDKI (D1-3) and relevant runs of relativistic reconnection (R1-3) are presented. The temperature (T/mc^2), the system size (L_x, L_y, L_z in unit of λ), the maximum non-thermal ratio of the kinetic energy (K_{nonth}/K), the energy conversion rate ($\tau_c d \ln(\Delta K)/dt$), the maximum energy (ε_{max}/mc^2) and their estimated values (ε_{est}/mc^2) are presented.